


BEST AVAILABLE COPY



Rec'd PCT/PTO 04 APR 2005
PCT/EP03/11725



#2

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


EP03/11725

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

P. Mahoney

Dated 4 November 2003

Patents Form 1/77 THE PATENT OFFICE

Patents Act 1977
(Rule 16)

29 OCT 2002

RECEIVED BY FAX

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

PPD 70159/GB/P

29OCT02 E759374-1 D02093
P01/7700 0.00-0225129.6

2. Patent application number

(The Patent Office will fill in this part)

0225129.6

29 OCT 2002

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

SYNGENTA Participations AG
Schwarzwaldallee 215
CH-4058 Basel
SWITZERLAND

Patents ADP number (*if you know it*)

8290835001

If the applicant is a corporate body, give the country/state of its incorporation

SWITZERLAND

4. Title of the invention

IMPROVEMENTS IN OR RELATING TO ORGANIC
COMPOUNDS

5. Name of your agent (*if you have one*)

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

Intellectual Property Department
Syngenta Limited
Jealott's Hill International Research Centre
PO Box 3538
Bracknell, Berkshire, RG42 6YA
UNITED KINGDOM

Patents ADP number (*if you know it*)

8029563001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)

Date of filing
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES (b)

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form

Description

19 /

Claim(s)

05 /

Abstract

00

Drawing(s)

00

10. If you are also filing any of the following,
state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right
to grant of a patent (Patents Form 7/77)Request for preliminary examination
and search (Patents Form 9/77)Request for substantive examination
(Patents Form 10/77)Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.
Syngenta LimitedSignature
Authorised Signatory

Date

29/10/02

12. Name and daytime telephone number of
person to contact in the United Kingdom

Joanna Carmen CHANDLER = 01344 414365

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

PPD 70159

- 1 -

DUPLICATE

IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS

The present invention relates to genetic engineering of plants and in particular to an insect resistant transgenic cotton plant. It also relates to methods of detecting material derived from the plant.

Plant pests are a major factor in the loss of the world's important agricultural crops. About \$8 billion is lost every year in the U.S. alone due to infestations of plants by non-mammalian pests including insects. In addition to losses in field crops, insect pests are also a burden to vegetable and fruit growers, to producers of ornamental flowers, and to home gardeners.

Insect pests are mainly controlled by intensive applications of chemical pesticides, which are active through inhibition of insect growth, prevention of insect feeding or reproduction, or cause death. Good control of insect pests can thus be reached, but these chemicals can sometimes also affect other, beneficial insects. Another problem resulting from the wide use of chemical pesticides is the appearance of resistant insect varieties. This has been partially alleviated by various resistance management practices, but there is an increasing need for alternative pest control agents. Biological pest control agents, such as *Bacillus thuringiensis* strains expressing pesticidal toxins like δ -endotoxins, have also been applied to crop plants with satisfactory results, offering an alternative or compliment to chemical pesticides. The genes coding for some of these δ -endotoxins have been isolated and their expression in heterologous hosts has been shown to provide another tool for the control of economically important insect pests. In particular, the expression of insecticidal toxins such as *Bacillus thuringiensis* δ -endotoxins in transgenic plants, has provided efficient protection against selected insect pests, and transgenic plants expressing such toxins have been commercialized, allowing farmers to reduce applications of chemical insect control agents.

Recently, a new family of insecticidal proteins produced by *Bacillus* sp. during the vegetative stages of growth (vegetative insecticidal proteins (VIPs)) has been identified. U.S. Patents 5,877,012, 6,107,279, and 6,137,033 describe *vip3A* toxin genes isolated

PPD 70159

- 2 -

from *Bacillus* species. The VIP3A toxins possess insecticidal activity against a wide spectrum of lepidopteran insects including but not limited to fall armyworm, *Spodoptera frugiperda*, black cutworm, *Agrotis ipsilon*, sugarcane borer, *Diatraea saccharalis*, and lesser cornstalk borer, *Elasmopalpus lignosellus*, and when expressed in transgenic plants, for example cotton, confer protection on the plant from insect feeding damage.

The cotton family, genus *Gossypium*, a member of the Malvaceae, consists of 39 species, of which *Gossypium hirsutum* is the most commonly cultivated species. Three other species are also cultivated: *G. arboreum*, *G. barbadense*, and *G. herbaceum*. These cultivated species are grown primarily for the seed hairs that are made into textiles. Cotton is suitable as a textile fibre because the mature dry hairs twist in such a way that fine strong threads can be spun from them. Other products, such as cottonseed oil, cake, and cotton linters are byproducts of fibre production.

Damage to cotton crops by insect pests throughout the world results in a significant yield loss each year. Effective control of these pests to minimise yield loss is of great economic importance. Examples of insect pests of cotton include Beet armyworm (*Spodoptera exigua*), Boll weevil (*Anthonomus grandis grandis*), Cabbage looper (*Trichoplusia ni*), Clouded plant bug (*Neurocolpus nubilus*), Cotton aphid (*Aphis gossypii*), Cotton bollworm (*Heliothrips zea*), Cutworms (*Feltia subterranea*, *Peridroma saucia*, *Agrotis ipsilon*), European corn borer (*Ostrinia nubilalis*), Fall armyworm (*Spodoptera frugiperda*), Seedling thrips (*Frankliniella* spp.), Soybean looper (*Pseudoplusia includens*), Stink bugs (*Nezara viridula*, *Acrosternum hilare*, *Euschistus servus*), Tarnished plant bug (*Lygus lineolaris*), Tobacco budworm (*Heliothis virescens*) and Whiteflies (*Trialeurodes abutilonea*, *Bemisia tabaci*).

Transformation and regeneration of cotton plants is now a well-established procedure, typically based on *Agrobacterium tumefaciens* mediated transfer of foreign DNA into cotton plant parts and regeneration of said plant parts in tissue culture into fully fertile, transgenic cotton plants.

There exists a requirement to generate a cotton plant that is insect resistant so that yield loss through damage to cotton crops by insect pests is reduced. An insect resistant cotton

PPD 70159

- 3 -

plant could reduce the need to apply chemical pesticides, which may be detrimental to other, beneficial insects and the environment.

Therefore, the present invention relates to an insect resistant transgenic cotton event, designated COT102. It also relates to methods of detecting plant material derived therefrom. "COT102 event" in the context of this application refers to the original insecticidal transgenic cotton plant described herein. "Insecticidal" as used herein refers to any inhibitory effect on an insect, including but not limited to reduced feeding, retarded growth, reduced fecundity, paralysis or death. "Fecundity" comprises all aspects related to reproduction such as reproductive ability, reproductive frequency and number of offspring. Also embraced by this invention is any plant material derived from the COT102 event, including seeds.

The COT102 event exhibits a novel genotype comprising two expression cassettes. The first cassette comprises a suitable promoter for expression in plants operably linked to a gene that encodes a VIP3A insecticidal toxin, useful in controlling a wide spectrum of lepidopteran insect pests, and a suitable polyadenylation signal. Suitable promoters may be isolated from, *inter alia*, plants. Numerous plant promoters have been isolated and characterised; one example of a suitable plant promoter is the Actin promoter (Act2) from rice. Additional elements such as enhancer sequences may also be incorporated into the expression cassette in order to boost levels of gene expression. Alternatively it may be desirable to include a targeting sequence, for example, to direct transportation of the VIP3A toxin to a particular cellular compartment. Numerous polyadenylation signals have been isolated and characterised. An example of a suitable polyadenylation signal functional in plants is that from the nopaline synthase gene (nos) of *Agrobacterium tumefaciens*.

The second cassette comprises a gene which, when expressed, can be used as a selectable marker. Numerous selectable markers have been characterised, including some that confer tolerance to antibiotics and others that confer tolerance to herbicides. An example of a suitable selectable marker gene is one that confers tolerance to a hygromycin. Further expression cassettes are optionally comprised in the COT102 event. For example these may provide other desirable benefits such as herbicide resistance.

PPD 70159

- 4 -

Thus according to the first aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1. In one embodiment said polynucleotide comprises at least 18 contiguous
5 nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises at least 20 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment still said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 1. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 1. In a preferred embodiment of the present invention,
10 there is provided a polynucleotide comprising the sequence of SEQ ID NO: 1.

In a further aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2. In one embodiment said polynucleotide comprises at least 18 contiguous nucleotides from SEQ
15 ID NO: 2. In a further embodiment said polynucleotide comprises at least 20 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment still said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 2. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 2. In a preferred embodiment of the present invention, there is provided a
20 polynucleotide comprising the sequence of SEQ ID NO: 2.

In another aspect of the present invention there is provided a plant comprising a polynucleotide which comprises at least 17 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. Preferably said plant comprises at least 18 contiguous nucleotides of
25 SEQ ID NO: 1 and / or SEQ ID NO: 2. More preferably said plant comprises at least 20 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. More preferably said plant comprises at least 22 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. More preferably still said plant comprises at least 24 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. Most preferably said plant comprises the
30 sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2. In an embodiment of the present invention, said plant is a cotton plant. Preferably, said plant is an insecticidal cotton plant which is the COT102 event, or a plant derived therefrom.

PPD 70159

- 5 -

The skilled man is familiar with plant transformation methods. Two principal techniques have been characterised across a wide range of plant species: transformation by *Agrobacterium* and transformation by direct DNA transfer.

- 5 *Agrobacterium*-mediated transformation is a preferred method for transformation of dicotyledonous plants. The foreign DNA to be introduced into the plant is cloned into a binary vector in between left and right border consensus sequences. This is the T-DNA region. The vector is transferred into an *Agrobacterium* cell, which is subsequently used to infect plant tissue. The T-DNA region of the vector comprising the foreign DNA is
10 inserted into the plant genome.

- Alternatively, direct DNA transfer can be used to introduce the DNA directly into a plant cell. One suitable method of direct transfer may be bombardment of plant cells with a vector comprising the DNA for insertion using a particle gun; another established
15 method, 'whiskers', involves coating the DNA onto silicon carbide fibres onto which cells are impaled.

- Following transformation, transgenic plants must be regenerated from the transformed plant tissue, and progeny possessing the foreign DNA selected using an appropriate
20 marker such as resistance to hygromycin.

- A plant of this aspect of the invention, as described herein, has an insecticidal effect on insects from one or more species from the group comprising *Heliothis* sp., *Helicoverpa* sp. and *Spodoptera* sp. which may infest it. "Infest" as used herein refers to attack,
25 feeding or damage in any way by one or more insects. Thus, for example, the plant of the present invention will provide a self-defence mechanism against infestation by pest insects such as *Helicoverpa zea* (cotton boll worm). As a result, a reduced number of insecticide sprays are required during the cultivation of said plant compared to a non-transgenic cotton plant of the same variety and yield loss through insect pests is kept at a
30 minimal level.

The present invention is not limited to the COT102 event itself, but is further extended to include any plant material derived therefrom, including seeds in so far as they contain

PPD 70159

- 6 -

one of the present inventive polynucleotides. The present invention includes, but is not limited to plants that are derived from a breeding cross with the COT102 event or a derivative therefrom by conventional breeding or other methods. The invention also includes plant material derived from the COT102 event that may comprise additional, modified or fewer polynucleotide sequences compared to the COT102 event or exhibit other phenotypic characteristics. For example it may be desirable to transform the plant material derived from the COT102 event to generate a new event that possess an additional trait, such as a second insect resistance gene which produces a toxin with a different mode of action or for the control of different insect species, or herbicide resistance. This process is known as gene stacking.

According to yet a further aspect of the present invention, there is provided a method of detecting plant material derived from the COT102 transgenic event comprising obtaining a sample for analysis; providing a pair of primers designed to bind to a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2; amplifying the region which lies between the sites at which the primers bind; and detecting the presence of the amplification product. Suitable pairs of primers for use in this method of detection can be designed using parameters well known to those skilled in the art of molecular biology now that SEQ ID NOs 1 and 2 are made available. The preferred sequence of said primers is included in the sequence listing as SEQ ID NO: 3 and SEQ ID NO: 4.

In an embodiment of the present invention, the region amplified by said method (the 'amplicon') is preferably between 300 and 1000 base pairs in length. More preferably the amplicon is between 500 and 900 base pairs in length. More preferably still the amplicon is 800 base pairs in length. Most preferably the amplicon is produced using the above method in conjunction with the primers of the sequence of SEQ ID NO: 3 and SEQ ID NO: 4, and is 800 base pairs in length.

There are many amplification methods that may be used in accordance with this aspect of the invention. The underlying principle, a known technique to those skilled in the art, is the polymerase chain reaction (PCR). The amplification product from a PCR reaction

PPD 70159

- 7 -

may be visualised by staining with ethidium bromide and excitation with UV light, typically after size separation using agarose gel electrophoresis.

An embodiment of the present invention employs variations of the PCR principle such as
5 TaqMan™. This involves labelling at least one of the primers involved in the amplification process with a fluorescent dye. When unbound, the primer adopts a conformation such that no fluorescence can be detected. However, when the primer is bound to a piece of DNA, the conformation changes and fluorescence can be detected. In this way, the amplification process can be monitored in real-time, the intensity of
10 fluorescence corresponding directly to the level of amplification. Further embodiments of the present invention include, but are not limited to, RACE PCR.

In another aspect of the invention there is provided a method of detecting plant material derived from the COT102 event comprising obtaining a sample for analysis; providing a
15 probe designed to bind to a polynucleotide which comprises at least 14 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2 when said polynucleotide is single stranded; hybridising said probe with the sample; and detecting whether the probe has hybridised. Preferably said probe comprises the sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2. In an embodiment of the present invention there is provided a method of
20 detecting plant material derived from the COT102 event using a probe selected from the group comprising SEQ ID NO: 5, SEQ ID NO: 6 and SEQ ID NO: 7. Preferably said probe comprises SEQ ID NO: 5. The probe may be, for example, a PCR product or restriction digestion fragment. In a further embodiment, the probe as described herein may be tagged with a fluorescent, radioactive, enzymatic or other suitable label to enable
25 hybridisation to be detected. The skilled man will know how to design suitable probes, now that he has the benefit of the present disclosure.

In a further embodiment of the present invention, there is provided a method of hybridising a probe to the sample under stringent conditions and detecting whether the
30 probe has hybridised. Stringent hybridisation conditions are well known to the skilled man and comprise, for example: hybridisation at a temperature of about 65°C in a solution containing 6 x SSC, 0.01% SDS and 0.25% skimmed milk powder, followed by rinsing at the same temperature in a solution containing 0.2 x SSC and 0.1% SDS.

PPD 70159

- 8 -

Suitable techniques for detecting plant material derived from the COT102 event based on the hybridisation principle include, but are not limited to Southern Blots, Northern Blots and In-Situ hybridisation. The skilled man is familiar with techniques such as these.

- 5 Typically, they involve incubating a probe with a sample, washing to remove unbound probe, and detecting whether the probe has hybridised. Said detection method is dependent on the type of tag attached to the probe – for example, a radioactively labelled probe can be detected by exposure to and development of x-ray film. Alternatively, an enzymatically labelled probe may be detected by conversion of a substrate to effect a
- 10 colour change.

- In a further aspect of the invention there is provided a method of detecting plant material derived from the COT102 event comprising obtaining a sample for analysis; providing an antibody designed to bind to a VIP protein contained within a plant comprising at least
- 15 14 contiguous nucleotides from SEQ ID NO: 1 and / or SEQ ID NO: 2; incubating said antibody with the sample; and detecting whether the antibody has bound. In one embodiment of the present invention said VIP protein comprises the sequence of SEQ ID NO: 8.

- 20 Suitable methods of detecting plant material derived from the COT102 event based on said antibody binding include, but are not limited to Western Blots, Enzyme-Linked ImmunoSorbent Assays (ELISA) and SELDI mass spectrometry. The skilled man is familiar with these immunological techniques. Typical steps include incubating a sample with an antibody that binds to the VIP protein, washing to remove unbound antibody, and
- 25 detecting whether the antibody has bound. Many such detection methods are based on enzymatic reactions - for example the antibody may be tagged with an enzyme and on application of a suitable substrate, a colour change detected. Suitable antibodies may be monoclonal or polyclonal.

- 30 In a further aspect of the present invention there is provided a method of detecting plant material derived from the COT102 event comprising obtaining a sample for analysis; subjecting one or more insects of the species *Spodoptera frugiperda* (susceptible to VIP3A) to the sample; subjecting one or more insects of species *Ostrinia nubilalis* (not

PPD 70159

- 9 -

susceptible to VIP3A) to the sample as a control; detecting whether the sample has an insecticidal effect on insects from each species; and comparing the results with an authentic COT102 bioassay profile. The results are compared against an authentic COT102 bioassay profile that is produced using insects of the same condition which have
5 been subjected to the same dose and type of COT102 plant material and where the insecticidal effect is detected the same length of time after subjecting the insects to the COT102 sample. *Spodoptera frugiperda* is a positive control for COT102 as it is susceptible to a suitable dose of VIP3A, while *Ostrinia nubilalis* is a negative control for COT102 as it is not susceptible to a suitable dose of VIP3A.

10

In one embodiment of the invention, the method of detecting plant material derived from the COT102 event includes but is not limited to leaf-feeding bioassays in which a leaf or other suitable plant part from the COT102 event or any plant material derived from the COT102 event, is infested with one or more pest insects. Detection may be through
15 assessment of damage to the leaf or plant part after set time periods, or assessment of mortality or another insecticidal effect on the insects.

In another aspect of the invention, there is provided a kit of parts comprising a means for detecting the presence of plant material derived from the COT102 event. Preferably, said
20 kit of parts comprises a means for detecting the presence of a polynucleotide comprising at least 14 contiguous nucleotides from the sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2, or a VIP protein. In an embodiment of the present invention, said kit of parts may comprise DNA amplification-detection technology such as PCR or TaqMan™. In a further embodiment of the present invention, said kit of parts may comprise probe
25 hybridisation-detection technology such as Southern Blots, Northern Blots or In-Situ Hybridisation. In another embodiment of the present invention, said kit of parts may comprise antibody binding-detection technology such as Western Blots, ELISA's or SELDI mass spectrometry. In a further embodiment of the present invention, said kit of parts may comprise insect bioassay-detection technology such as leaf feeding bioassays
30 or mortality bioassays. In a further embodiment of the present invention, said kit of parts may comprise any combination of the afore-mentioned detection technologies.

PPD 70159

- 10 -

EXAMPLES

The invention will be further apparent from the following non-limiting examples in conjunction with the associated sequence listings as described below:

5

SEQ ID NO 1: Polynucleotide sequence which extends across the junction where the 5' end of the expression cassette is inserted into the cotton genome in event COT102.

10

SEQ ID NO 2: Polynucleotide sequence which extends across the junction where the 3' end of the expression cassette is inserted into the cotton genome in event COT102.

SEQ ID NO 3: Polynucleotide sequence suitable for use as a primer in the detection of the COT102 event.

15

SEQ ID NO 4: Polynucleotide sequence suitable for use as a primer in the detection of the COT102 event.

SEQ ID NO 5: Polynucleotide sequence suitable for use as a probe in the detection of the COT102 event.

SEQ ID NO 6: Polynucleotide sequence suitable for use as a probe in the detection of the COT102 event.

20

SEQ ID NO 7: Polynucleotide sequence suitable for use as a probe in the detection of the COT102 event.

SEQ ID NO 8: Amino acid sequence of the VIP3Aa toxin protein.

25 **Example 1: Cloning and Transformation**

1.1 Vector cloning

Standard gene cloning techniques of restriction digestion and ligation of fragments from in-house vectors were used to construct the transformation vector, pNOV3001. The
30 vector included a selection cassette comprising a Ubiquitin (UBQ3) promoter, the UBQ3 intron, a gene sequence which encodes a protein conferring resistance to hygromycin, and a nos polyadenylation sequence. The vector also included the expression cassette of the target gene, which cassette comprised an Actin (Act2) promoter, the Act2 intron, a

PPD 70159

- 11 -

sequence encoding the VIP3Aa gene that had been codon optimised for expression in maize, and a nos polyadenylation sequence. The vector also comprised a gene conferring resistance to an antibiotic, spectinomycin, for prokaryotic selection.

- 5 The vector was transformed into *Agrobacterium tumefaciens* strain EHA101 using standard *Agrobacterium* transformation techniques, and transformed cells selected through their resistance to spectinomycin.

1.2 Plant transformation

- 10 The COT102 event was produced by *Agrobacterium*-mediated transformation of *Gossypium hirsutum* L. cv Coker 312.

Coker 312-17 seeds were surface sterilized for 30 seconds in 70% ethanol using enough ethanol to cover the amount of seed to be sterilized. The seeds were washed with
15 ethanol, rinsed in sterile water and soaked in a 12% Clorox + Tween 20 solution for 20 minutes. This washing procedure was performed 3 times. Seeds were then placed onto germination medium (Stewart and Hsu, 1977) and allowed to germinate at 30°C for 7 – 10 days.

- 20 2 ml cultures of *Agrobacterium* containing the pNOV3001 construct were grown overnight in appropriate antibiotics and then diluted with MSNH medium (19:1) in a sterile petri dish. Hypocotyls were cut into 6 – 8 mm lengths and placed in the diluted *Agrobacterium* solution for at least 30 seconds. Hypocotyl explants were removed from the *Agrobacterium* solution and blotted on sterile filter paper to remove excess bacteria.
25 Hypocotyls were placed onto T2 medium (MS salts, B5 vitamins, 0.1 mg/L 2, 4-D, 0.5 mg/L kinetin, 30 g/L glucose, 2 g/L Phytigel – pH 5.8) and co-cultured with the *Agrobacterium* for 72 hours in the dark.

- Hypocotyl explants were again blotted on sterile filter paper and transferred to plates
30 containing MS2NK medium (MS salts, B5 vitamins, 2 mg/L NAA, 0.1 mg/L kinetin, 30 g/L glucose, 2 g/L Phytogel, 500 mg/L cefotaxime, 10 mg/L hygromycin – pH 5.8). The plates were wrapped with parafilm and incubated in the light at 30°C for several months until callus was formed.

PPD 70159

- 12 -

Callus was broken up as small as possible and placed in a 50 ml Erlenmeyer flask containing 10 ml of liquid MSNH medium (MS salts, B5 vitamins, 30 g/L glucose – pH 5.8). The suspended callus was shaken at 110 rpm in the light at 30°C until small white
5 slightly round cell clusters were visible. Cells were washed and plated onto solid MSNH medium (MS salts, B5 vitamins, 30 g/L glucose, 2 g/L Phytogel – pH 5.8). Plates were checked monthly for somatic embryo development.

Mature somatic embryos were picked from the plates and placed onto plates containing
10 SA medium (Stewart and Hsu salts, 20 g/L sucrose, 20 g/L agar – pH 5.8). The plates of embryos were placed in the dark for approximately 14 days. Roots were trimmed from the maturing embryos and the embryos were transferred to SGA medium (Stewart and Hsu salts, 5 g/L sucrose, 1.5 g/L Phytogel, 5 g/L agar – pH 6.8).

15 After the first true leaf emerged, young plants were moved to pint sized canning jars containing SGA medium. When the plants reached 7 – 10 cm in height, the top was cut off and transferred to another jar. On developing a good root system, the thus rooted cuttings were transplanted into pots and grown in the greenhouse.

20 1.3 Identification and selection of transgenics

Putative transgenic plants were screened by PCR for the presence of the VIP3A gene. Positive events were identified and screened using insect bioassays for insecticidal activity against Fall Armyworm (*Spodoptera frugiperda*) (see Example 5). Insecticidal lines were characterized for copy number by TaqMan™ analysis (see Example 2). T1
25 seed from 3 single copy & 2 double copy events were observed in a field trial for insect resistance and agronomic quality. Two events, COT101 and COT102, were chosen based on having a single copy of the transgene, good protein expression as identified by ELISA (see Example 4), good insecticidal activity against Cotton Boll Worm (*Helicoverpa zea*) and field performance. At the end of the second year of field trials,
30 results between the two events were compared and COT102 was progressed.

PPD 70159

- 13 -

1.4 Verification of sequence of COT102

Genomic DNA was isolated from the COT102 event. This was used in the sequencing of the junctions of the DNA insertion site with the cotton genomic DNA in the COT102 event, using standard DNA sequencing techniques.

5

Example 2: COT102 Detection via TaqMan™

DNA was extracted from leaf tissue using the Wizard™ Magnetic 96 DNA Plant System (Promega, #FF3760), according to the manufacturers instructions, with an additional step at the beginning of the protocol: following grinding of the leaf material, 0.9ml Cotton Extraction Buffer (0.2M Tris pH 8.0, 50mM EDTA, 0.25M NaCl, 0.1% v/v 2-mercaptoethanol, 2.5% w/v polyvinyl-pyrrolidone) was added to each well, the plant tissue resuspended and the plate centrifuged at 4,000 rpm (2755g) for 10 minutes. After aspirating and discarding the supernatant, 300ul Lysis Buffer A (Promega) was added and the manufacturers protocol was followed from this point. This procedure resulted in approximately 85ul of purified genomic DNA at a concentration of approximately 10ng/ul.

20 TaqMan™ PCR reactions were setup using a standard reaction mix comprising:

- 625ul 2x Jumpstart Master Mix for Q-PCR (Sigma, #P2893), supplemented with 15mM MgCl₂ and 200nM Strata-ROX
- 25ul 50x FAM primer / probe mix
- 25ul 50x TET primer / probe mix
- 25 200ul Water.

50x primer / probe mixes comprise 45ul of each primer at a concentration of 1mM, 50ul of the probe at a concentration of 100uM and 860ul 1x TE, and were stored in an amber tube at 4°C.

30

Examples of suitable primer / probe sequence combinations which were used are:

PPD 70159

- 14 -

<u>Primer Name</u>	<u>Primer Sequence 5'-3'</u>	
GhCHI2b-F	GGTCCCTGGATACGGTGTCA	Forward
GhCHI2b-R	TTGAGGGTTGGATCCTTTGC	Reverse
GhCHI2b-TET	CCAACATCATCAATGGTGGCATCGAAT	Probe
	(5' label = TET, 3' label = TAMRA)	
Hygromycin-F	CAGGCAGGTCTTGCAACGT	Forward
Hygromycin-R	CGAGAGCCTGACCTATTGCAT	Reverse
Hygromycin-FAM	ACACCCTGTGCACGGCGGG	Probe
	(5' label = FAM, 3' label = TAMRA)	
Vip3-F	ATGAAGACCCTGCGCTACGA	Forward
Vip3-R	ACGCCCAGTGGCATGTAGA	Reverse
Vip3-FAM	AGCGAGGCCGAGTACCGCACC	Probe
	(5' label = FAM, 3' label = TAMRA)	

- 7ul of master mix was dispensed into each well of a 384-well TaqMan™ assay plate. 3ul DNA template was added to the appropriate wells. 3ul of copy control dilution series was added to specific wells as a control. The reactions are run in an ABI7900 (Applied Biosystems) using the following cycling conditions:

	Step	Temperature	Time
	1	50°C	2 min
	2	95°C	10 min
10	3	95°C	15 sec
	4	60°C	1 min
	5	Goto step 3, repeat 40 times	

Data was analysed using SDS2.0 software (Applied Biosystems).

PPD 70159

- 15 -

Example 3: COT102 Detection via Southern Blot

DNA suitable for use in Southern Blotting was extracted using the following procedure.

5

Approximately 5 to 10 grams of plant tissue was ground in liquid nitrogen using a mortar and pestle. The plant tissue was resuspended in 12.5ml extraction buffer A (0.2M Tris pH8.0, 50mM EDTA, 0.25M NaCl, 0.1% v/v B-mercaptoethanol, 2.5% w/v Polyvinyl-pyrrolidone), and centrifuged for 10 minutes at 4,000 rpm (2755g). After discarding the
10 supernatant the pellet was resuspended in 2.5ml Extraction Buffer B (0.2M Tris pH8.0, 50mM EDTA, 0.5M NaCl, 1% v/v B-mercaptoethanol, 2.5% w/v Polyvinyl-pyrrolidone, 3% Sarkosyl, 20% Ethanol) and incubated at 37°C for 30 minutes. During incubation, the sample was mixed once with a sterile loop. After incubation an equal volume of chloroform/isoamyl alcohol (24:1) was added, mixed gently by inversion, and
15 centrifuged for 20 minutes at 4,000 rpm. The aqueous layer was collected, and 0.54 volume isopropanol added followed by centrifugation for 5 minutes at 4,000 rpm to precipitate the DNA. The supernatant was discarded and the DNA pellet resuspended in 500ul TE. In order to degrade any RNA present, the DNA was incubated at 37°C for 30 minutes with 1ul 30mg/ml RNase A, centrifuged for 5 minutes at 4,000 rpm and
20 precipitated by centrifugation at 14,000 rpm for 10 minutes in the presence of 0.5 volume 7.5M ammonium acetate and 0.54 volume isopropanol. After discarding the supernatant, the pellet was washed with 500ul 70% ethanol and allowed to dry before resuspending in 100ul TE.

25 The DNA was quantified using a spectrophotometer or fluorometer (using 1xTNE and Hoechst dye). Suitable enzyme digests were prepared using 8ug DNA per digest in a total volume of 50ul. Digests included BamHI, EcoRI, EcoRV, HindIII, NcoI, SacI, ScaI, SpeI and PstI, both alone and in combination. Digests were incubated overnight at the appropriate temperature for each enzyme. Samples were spun in a speed vacuum to
30 reduce the volume to 30ul. Bromophenol blue loading dye was added and each sample loaded on a 0.8% TBE agarose gel containing ethidium bromide. The gel was run at 60 volts overnight.

PPD 70159

- 16 -

The gel was washed in 0.25M HCl for 15 minutes to depurinate the DNA, and then washed with water. A Southern Blot was set up as follows: 20 sheets of thick dry blotting paper was placed in a tray and 4 sheets of thin dry blotting paper placed on top. One sheet of thin blotting paper was pre-wet in 0.4M NaOH and placed on top of the stack, followed by a sheet of Hybond-N+ transfer membrane (Amersham Pharmacia Biotech, #RPN303B), also pre-wet in 0.4M NaOH. The gel was placed on top ensuring that there were no air bubbles between the gel and the membrane. Three further sheets of pre-soaked blotting paper were placed on top of the gel and the buffer tray filled with 0.4M NaOH. Connecting the gel stack with the buffer tray using a wick pre-soaked in 0.4M NaOH started the transfer of DNA to the membrane. DNA transfer took place for approximately 4 hours at room temperature. Following transfer, the Hybond membrane was rinsed in 2x SSC for 10 seconds and the DNA bound to the membrane via UV cross-linking.

A suitable DNA probe was prepared by PCR. 25ng probe DNA in 45ul TE was boiled for 5 minutes, placed on ice for 7 minutes then transferred to a Rediprime II (Amersham Pharmacia Biotech, #RPN1633) tube. After addition of 5ul P32-labelled dCTP to the Rediprime tube, the probe was incubated at 37°C for 15 minutes. The probe was cleaned by centrifugation through a microspin G-50 column (Amersham Pharmacia Biotech, #27-5330-01) according to the manufacturers instructions. The activity of the probe was measured using a scintillation counter.

The Hybond membrane was pre-hybridised by wetting with 20ml pre-warmed Church pre-hybridisation solution (500mM NaPO₄, 1mM EDTA, 7% SDS, 1% BSA) at 65°C for 30 minutes. The labelled probe was boiled for 5 minutes, and placed on ice for 10 minutes. An appropriate amount of probe (1 million counts per 1ml pre-hybridisation buffer) was added to the pre-hybridisation buffer and hybridisation occurred at 65°C overnight. The following day, the hybridisation buffer was discarded, and following a rinse with 20ml Church Wash Solution 1 (40mM NaPO₄, 1mM EDTA, 5% SDS, 0.5% BSA), the membrane washed in 150ml Church Wash Solution 1 at 65°C for 20 minutes. This process was repeated twice with Church Wash Solution 2 (40mM NaPO₄, 1mM EDTA, 1% SDS). The membrane was exposed to a phosphor screen or X-ray film to detect where the probe has bound.

PPD 70159

- 17 -

Example 4: COT102 Detection via ELISA

5 Cotton tissue for analysis was harvested and frozen at -70°C. Fresh tissue was ground to a fine powder and weighed into a labelled polypropylene tube. Extraction buffer (100mM Tris, 100mM Sodium Borate, 5mM MgCl, 0.05% Tween 20, 0.2% Sodium Ascorbate, Water, pH 7.8, 1mM AEBSF, 0.001mM Leupeptin) was added to the sample in a ratio of 2:1 (volume extraction buffer : sample fresh weight) for fresh tissue or 30:1
10 (volume extraction buffer : sample dry weight) for lyophilised tissue. The sample was vortexed and homogenised using a Brinkman PT 10/35 Polytron equipped with a PTA 10TS foam-reducing generator, until the mixture became liquefied. Extracts were centrifuged at 10,000 x g for 15 minutes. The protein extract supernatant was stored at 2-8°C.

15

The ELISA procedure used standard techniques as follows. A 96-well plate was soaked in ethanol for 2 hours, and air-dried. The plate was coated with 50ul goat anti-VIP3A antibody per well and incubated overnight at 2-8°C. After washing three times with 1X ELISA wash solution (100mM Tris, 0.5% Tween-20, 75mM NaCl, pH8.5), the plate was
20 dried briefly by tapping upside down on a paper towel. 150ul blocking solution (10mM NaPO₄, 140mM NaCl, 1% BSA, 0.02% Sodium Azide, titrated to pH7.4 with monobasic NaPi and dibasic NaPi) was added to each well followed by incubation at room temperature for 45 minutes. The plate was washed 3 times as described above.

25 VIP3A standards and protein extract samples were applied to appropriate wells of the plate in triplicate, 50ul total volume per well. The plate was incubated at 2-8°C for 1 hour 30 minutes, followed by room temperature for a further 30 minutes. The plate was washed three times with ELISA wash solution, and then incubated at 35-39°C for 1 hour with 50ul rabbit anti-VIP3A antibody per well. The plate was washed three times with
30 ELISA wash solution, and incubated at room temperature for 30 minutes with 50ul donkey anti-rabbit alkaline phosphatase per well. Following a further three washes with ELISA wash solution, 50ul phosphatase substrate solution was added per well and the plate incubated for 30 minutes at room temperature. The reaction was stopped by

PPD 70159

- 18 -

addition of 50ul 3M NaOH per well. The absorbance of the solution in each well was measured at 405nm using a Ceres 900C multiwell plate reader and the results analysed using KC3 Curve fitting software (Bio-Tek Instruments Inc.). The concentration of VIP3A in the samples was calculated by reference to the VIP3A protein standards.

5

Example 5: COT102 Detection via Insect Bioassay

Leaf assays were performed on Fall Army Worm (*Spodoptera frugiperda*), Cotton Boll
10 Worm (*Helicoverpa zea*) and Tobacco Budworm (*Heliothis virescens*) as follows:
Pads were soaked with 300ul to 500ul distilled water and placed into Gelman dishes.
Leaf pieces measuring between approximately 0.5 square inches and 0.75 square inches
were excised from cotton plants 8 to 12 inches in height, and placed on the pads.
Between 8 and 10 insect larvae were placed in each dish and a lid fitted. The dishes were
15 incubated at 28°C. On the third and sixth days after infestation, damage to the leaf in
each dish was scored and compared with the control plants.

As an alternative, a 'boll' assay was performed as follows:

Four absorbent pads were saturated with water and placed inside a large plastic cup.
20 Three extra thick glass filters, each soaked with 100ul distilled water, were placed in a
smaller plastic cup, which was then seated inside the larger cup. A 1.25 inch long boll
was excised, immersed in 10mg/ml to 20mg/ml Nystatin and placed on the filters in the
small cup. 50 insect larvae were placed on the square or boll and a lid attached to the
larger cup. The squares or bolls were re-infested with 50 more larvae after 7 days.

25

The experiment was incubated at room temperature for approximately 3 weeks. The
bolls were then cut open to determine damage. Damage to the boll was compared to the
control samples.

PPD 70159

- 19 -

REFERENCES

U.S. Patent 5,877,012 : Estruch et al. (Novartis Finance Corporation)

5 U.S. Patent 6,107,279 : Estruch et al. (Novartis Finance Corporation)

U.S. Patent 6,137,033 : Estruch et al. (Novartis Finance Corporation)

Stewart JM & Hsu CL (1977) Planta 137 p.113117

PPD 70159

- 20 -

CLAIMS

1. A polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1.

5

SEQ ID 1: GGC AAA TAT TCA GGT AAA CAA ATT GA

2. A polynucleotide according to claim 1 comprising at least 18 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1.

10

3. A polynucleotide according to claim 1 comprising at least 20 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1.

4. A polynucleotide according to claim 1 comprising the sequence of SEQ ID NO: 1.

15

5. A polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2.

SEQ ID 2: CTA TCA GTG TTT AAT AAA TAT GGG CA

20

6. A polynucleotide according to claim 5 comprising at least 18 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2.

7. A polynucleotide according to claim 5 comprising at least 20 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2.

25

8. A polynucleotide according to claim 5 comprising the sequence of SEQ ID NO: 2.

9. An insect resistant plant comprising a VIP3A protein and a polynucleotide according to any of claims 1 to 8.

30

10. A plant according to claim 9 which is a cotton plant.

PPD 70159

- 21 -

11. An insecticidal cotton plant according to claim 10 which is derived from the COT102 event.

12. A method of detecting plant material derived from the COT102 event comprising:

- 5 • obtaining a sample for analysis;
- providing a pair of primers designed to bind to a polynucleotide as claimed in claims 1 to 8 when said polynucleotide is single stranded;
- amplifying the region which lies between the sites at which the primers bind; and
- detecting the presence of the amplification product;
- 10 whereby the presence of the amplification product is indicative that the sample is derived from the COT102 event.

13. A method according to claim 12 wherein the first primer has the sequence of SEQ ID NO: 3 and the second primer has the sequence of SEQ ID NO: 4.

15

SEQ ID 3: 5' - AAGGACGTGAGCGAGATGTT -3'

SEQ ID 4: 5' - TGTGACACCGATCCACCTAA -3'

14. A method of detecting plant material derived from the COT102 event comprising:

- 20 • obtaining a sample for analysis;
- providing a probe designed to bind to a polynucleotide as claimed in claims 1 to 8 when said polynucleotide is single stranded;
- hybridising said probe with the sample; and
- detecting whether the probe has hybridised.
- 25 whereby the hybridisation of the probe is indicative that the sample is derived from the COT102 event.

15. A method according to claim 14 wherein the sequence of the probe is selected from the group comprising SEQ ID NO: 5, SEQ ID NO: 6 and SEQ ID NO: 7.

30

SEQ ID 5:

GACAAGGACA GCTTGAGCGA GGTGATCTAC GGCGACATGG ACAAGCTGCT
GTGTCCGGAC CAGAGCGAGC AAATCTACTA CACCAACAAC ATCGTGTTC

PPD 70159

- 22 -

CGAACGAGTA CGTGATCACC AAGATCGACT TCACCAAGAA GATGAAGACC
CTGCGCTACG AGGTGACCGC CAACTTCTAC GACAGCAGCA CCGGCGAGAT
CGACCTGAAC AAGAAGAAGG TGGAGAGCAG CGAGGCCGAG TACCGCACCC
TGAGCGCGAA CGACGACGGC GTCTACATGC CACTGGGCGT

5

SEQ ID 6:

AGGTAGTTGG CGTCATCGAG CGCCATCTCG AACCGACGTT GCTGGCCGTA
CATTTGTACG GCTCCGCAGT GGATGGCCGC CTGAAGCCAC ACAGTGATAT
TGATTTGCTG GTTACGGTGA CCGTAAGGCT TGATGAAACA ACGCGGCGAG
10 CTTTGATCAA CGACCTTTTG GAAACTTCGG CTTCCCCTGG AGAGAGCGAG
ATTCTCCGCG CTGTAGAAGT CACCATTGTT GTGCACGACG ACATCATTCG
GTGGCGTTAT CCAGCTAAGC GCGAACTGCA ATTTGGAGAA TGGCAGCGCA
ATGACATTCT TGCAGGTATC TTCGAGCCAG CCACGATCGA CATTGATCTG
GCTATCTTGC TGACAAAAGC AAGAGAACAT AGCGTTGCCT TGGTAGGTCC

15 AG

SEQ ID 7:

CGCCGTGCAC AGGGTGTAC GTTGCAAGAC CTGCCTGAAA CCGAACTGCC
CGCTGTTCTG CAGCCGGTCG CGGAGGCCAT GGATGCGATC GCTGCGGCCG
20 ATCTTAGCCA GACGAGCGGG TTCGGCCCAT TCGGACCGCA AGGAATCGGT
CAATACACTA CATGGCGTGA TTTCATATGC GCGATTGCTG ATCCCCATGT
GTATCACTGG CAAACTGTGA TGGACGACAC CGTCAGTGCG TCCGTGCGCG
AGGCTCTCGA TGAGCTGATG CTTTGGGCCG AGGACTGCCC CGAAGTCCGG
CACCTCGTGC ACGCGGATTT CGGCTCCAAC AATGTCCTGA CGGACAAAT

25

16. A method according to claims 14 or 15 wherein the probe hybridises to the sample under stringent hybridisation conditions.

17. A method of detecting plant material derived from the COT102 event comprising:

- 30 • obtaining a sample for analysis;
- providing an antibody designed to bind to a VIP protein contained within a plant according to claims 9 to 11;
- incubating said antibody with the sample; and
- detecting whether the antibody has bound.
- 35 whereby the presence of antibody which has bound is indicative that the sample is derived from the COT102 event.

PPD 70159

- 23 -

18. A method according to claim 17 wherein the VIP protein has the sequence of SEQ ID NO: 8.

5 SEQ ID 8:

1 MNKNNTKLST RALPSFIDYF NGIYGFATGI KDIMNMIFKT
41 DTGGDLTLDE ILKNQQLLND ISGKLDGVNG SLNDLIAQGN
81 LNTELSKEIL KIANEQNOVL NDVNNKLDAL NTMLRVYLPK
121 ITSMLSDVMK QNYALSLOIE YLSKQLQEIS DKLDIINVNV
10 161 LINSTLTEIT PAYQRIKYVN EKFEELTFAT ETSSKVKKDG
201 SPADILDELT ELTELAKSVT KNDVDGFEFY LNTFHDVMVG
241 NNLFGRSALK TASELITKEN VKTSGSEVGN VYNFLIVLTA
281 LQAQAFLLT TCRKLLGLAD IDYTSIMNEH LNKEKEEFRV
321 NILPTLSNTF SNPNYAKVKG SDEDAKMIVE AKPGHALIGF
15 361 EISNDSITVL KVEAKLKQN YQVDKDSLSE VIYGDMDKLL
401 CPDQSEQIYY TNNIVFPNEY VITKIDFTKK MKTLRYEVTA
441 NFYDSSTGEI DLNKKKVESS EAEYRTLSAN DDGVYMPFGV
481 ISETFLTPIN GFGLQADENS RLITLTCKSY LRELLLATDL
521 SNKETKLIVP PSGFISNIVE NGSIEEDNLE FWKANNKNAY
20 561 VDHTGGVNGT KALYVHKDGG ISQFIGDKLK PKTEYVIQYT
601 VKGKPSIHLK DENTGYIHYE DTNNNLEDYQ TINKRFTTGT
641 DLKGVYLILK SQNGDEAWGD NFIIIEISPS EKLLSPELIN
681 TNNWTSTGST NISGNTLTLY QGGRGILKQN LQLDSPSTYR
721 VYFSVSGDAN VRIRNSREVL FEKRYMSGAK DVSEMFTTKF
25 761 EKDNFYIELS QGNNLYGGPI VHFYDVSIK

19. A method of detecting plant material derived from the COT102 event comprising:

- obtaining a sample for analysis;
- subjecting one or more insects of the species *Spodoptera frugiperda* to the sample;
- subjecting one or more insects of species *Ostrinia nubilalis* to the sample as a control;

PPD 70159

- 24 -

- detecting whether the sample has an insecticidal effect on insects from each species; and
- comparing the results with an authentic COT102 bioassay profile.

5 20. A kit of parts comprising a means for detecting the presence of plant material derived from the COT102 event.

21. A kit of parts according to claim 20 comprising a means for detecting the presence of a polynucleotide according to claims 1 to 8 or a VIP protein.

10

22. A kit of parts according to claims 20 to 21 comprising one or more of the methods according to claims 12 to 19.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.